arXiv:1303.5587v1 [cond-mat.str-el] 22 Mar 2013

Topological and symmetry broken phases of $\mathbb{Z}_N$ parafermions in one dimension

Roberto Bondesan and Thomas Quella

Institute of Theoretical Physics, University of Cologne
Zülpicher Straße 77, D-50937 Cologne, Germany

Emails: Roberto.Bondesan@uni-koeln.de, Thomas.Quella@uni-koeln.de

Abstract

We classify the gapped phases of $\mathbb{Z}_N$ parafermions in one dimension and construct a representative of each phase. Even in the absence of additional symmetries besides parafermionic parity, parafermions may be realized in a variety of phases, one for each divisor n of N. The phases can be characterized by spontaneous symmetry breaking, topology, or a mixture of the two. Purely topological phases arise if n is a unitary divisor, i.e. if n and N/n are co-prime. Our analysis is based on the explicit realization of all symmetry broken gapped phases in the dual $\mathbb{Z}_N$-invariant quantum spin chains.

1 Introduction

Topological phases of matter received a lot of interest recently due to their peculiar properties such as the existence of robust edge modes, non-abelian excitations and their potential applications to quantum computation [1, 2]. While one-dimensional bosonic systems do not exhibit intrinsic topological order [3], the situation is very different for fermions. According to the general theory, there should exist two phases of fermions which are described by a $\mathbb{Z}_2$ topological invariant [4, 5, 6]. An explicit realization of both phases is provided by Kitaev's Majorana chain which models the dynamics of spinless fermions in a quantum wire, in proximity with a p-wave superconductor [7].

The most remarkable feature of Kitaev's model is the emergence of stable gapless Majorana edge modes in the topologically non-trivial phase [7]. In view of their non-abelian statistics, being able to create and to manipulate such isolated Majorana fermions would be a major step towards the realization of topological quantum computers [8]. While Kitaev's chain merely serves as a proof of principle, by now more realistic experimental setups for the creation and detection of Majorana fermions have been suggested [9, 10]. The basic ingredients are spin orbit quantum wires in proximity with an s-wave superconductor and in a magnetic field. Recent experiments based on this setup provide first evidence that Majorana fermions indeed exist [11, 12] even though at least part of the observed features could also be explained by disorder effects [13, 14, 15].

More recently, it has been investigated whether similar phenomena can be realized in heterostructures involving *fractional* topological insulators. It was found that such systems can be used to isolate parafermions which may be regarded as fractionalized Majorana fermions [16, 17]. These efforts received their motivation from notes of P. Fendley on the existence of edge modes in parafermionic chains, unpublished for a long time but eventually made available in Ref. [18]. Other work on this subject includes Refs. [19, 20].

The study of parafermions has a long history. The exploration of para-particles began in Refs. [21, 22] in an attempt to generalize the canonical commutation relations of bosons and fermions. In contrast, these days the predominant notion of parafermions arises in the context of statistical physics, in connection with the so-called $\mathbb{Z}_N$ clock model [23]. This model generalizes the $\mathbb{Z}_2$ symmetry of the Ising model in a natural way to the $\mathbb{Z}_N$ symmetry of a clock with N marks.[1]

Just as the Ising model may be mapped to a problem of free fermions, using a variation of the non-local Jordan-Wigner transformation, the $\mathbb{Z}_N$ clock model exhibits a dual description in terms of fundamental parafermionic degrees of freedom. The parafermionic operators satisfy a $\mathbb{Z}_N$-graded variant of the usual Clifford algebra relations. This duality generalizes the well-known duality between the Ising model and free fermions for $N = 2$. It is known that the topologically trivial and the non-trivial phase of the Majorana chain are equivalent to the ordered and the disordered phase of the Ising model, respectively [26].

The previous paragraphs suggest two natural avenues to the classification of parafermionic chains. One first of all can generalize the group theoretical approach of Ref. [6] that has been applied successfully to enumerate and characterize the different phases of Majorana chains with or without time-reversal symmetry. This approach has the advantage of yielding the maximal set of *possibilities* in a clean and conceptually clear way. On the other hand, the proof that all topological phases can actually be realized requires a method of combining phases such that their topological charges add up. This procedure might not necessarily lead to the most natural representative of the new phase. Moreover, the approach is limited to phases without broken symmetry. This approach has been followed recently in Ref. [36].

The second approach which shall be followed in this paper is a complete characterization of gapped symmetric and symmetry broken phases of $\mathbb{Z}_N$ quantum spin chains. These can be engineered systematically on the level of ground states and their associated frustration free parent Hamiltonians [27]. Via the duality mapping mentioned before, this classification provides an explicit characterization of all parafermionic phases which cannot be connected without crossing a phase transition, i.e. closing the gap.

We shall find that the distinct parafermionic phases are labeled by the possible divisors of N. Generically they exhibit coexistence of both ferromagnetic and topological order. Only part of the degeneracy of the ground state of an open chain is topologically protected and due to gapless edge modes, and the rest can be lifted by spontaneous symmetry breaking. When the chain is closed, only the topological degeneracy disappears, leaving in general with a degenerate ground state. In particular, for a given divisor n of N, the phase will be purely topological if N and N/n are co-prime, and will exhibit a unique ground state in a closed system and gapless edge modes in an open system. If instead $N/n = \gcd(n, N/n)$ the phase is topologically trivial and entirely characterized by symmetry breaking. The details of the classification will be explained through concrete realizations of the phases in the text.

We would like to stress that our classification distinguishes only phases which cannot be connected by any local $\mathbb{Z}_N$-invariant interaction. Clearly, specific models with an explicit Hamiltonian depending on a given set of coupling constants may have arbitrarily complicated phase diagrams.

The paper is organized as follows. In Section 2 we introduce the kinematical setup that we are working with. This includes a detailed discussion of the symmetry algebra and of the Hilbert spaces in question. The possible symmetry breaking patterns and order parameters of the $\mathbb{Z}_N$ quantum spin chain are analyzed in Section 3 and a representative Hamiltonian is constructed for each phase. These results form the heart of our paper. Finally, we perform a non-local duality transformation in Section 4 and interpret our previous results from a parafermionic perspective. As it turns out, the original symmetry broken phases of the spin chains generally exhibit a mixture of symmetry breaking and topological protection. We end with concluding remarks and an outlook to open problems.

---

[1] The name "parafermions" is sometimes also used for the basic fields in a family of $\mathbb{Z}_N$ symmetric conformal field theories that arise at critical points of the clock models [24]. These parafermions may be used to construct the Read-Rezayi states, i.e. they are of relevance for the study of fractional quantum Hall systems [25]. In the present paper, we shall be concerned exclusively with the gapped phases of $\mathbb{Z}_N$ parafermions.

2 Parafermions and $\mathbb{Z}_N$ spin chains

We introduce the mathematical framework within which our considerations take place. This includes a detailed discussion of the symmetry algebra and of the Hilbert spaces involved.

2.1 $\mathbb{Z}_N$ spin chains

$\mathbb{Z}_N$ quantum spin chains are $\mathbb{Z}_N$ symmetric models where the spins can show in any of N directions in the plane which can be thought of as corresponding to the N distinct N^{th} roots of unity on the unit circle. This setup generalizes the familiar $\mathbb{Z}_2$ case of the Ising model where one deals with spins pointing either up or down, and at the same time, provides a specialization of the XY model in which the spins are restricted to take discrete values.

If we denote the size of the chain by L, the Hilbert space of the chain will be given by

$$\mathcal{H} = (\mathbb{C}^N)^{\otimes L} \,. \tag{2.1}$$

We choose a basis in $\mathcal{H}$ given by $|q_1, \dots, q_L\rangle$, $q_i \in \mathbb{Z}_N$. (This means that $|q\rangle \equiv |q+N\rangle$.) Define the operators

$$\sigma_i = \mathbf{1} \otimes \cdots \otimes \underbrace{\sigma}_{i} \otimes \mathbf{1} \cdots \otimes \mathbf{1} \tag{2.2}$$

$$\tau_i = \mathbf{1} \otimes \cdots \otimes \underbrace{\tau}_{i} \otimes \mathbf{1} \cdots \otimes \mathbf{1} \,, \tag{2.3}$$

where σ and τ are

$$\sigma = \begin{pmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & \omega & 0 & \dots & 0 \\ 0 & 0 & \omega^2 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & 0 & \omega^{N-1} \end{pmatrix} , \quad \tau = \begin{pmatrix} 0 & 1 & 0 & \dots & 0 \\ 0 & 0 & 1 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & \dots & 0 & 0 & 1 \\ 1 & \dots & 0 & 0 & 0 \end{pmatrix} . \tag{2.4}$$

σ, τ satisfy the relations:[2]

$$X^N = 1 \,, X^\dagger = X^{-1} \,, \quad \text{for } X = \sigma, \tau \tag{2.5}$$

$$\sigma^r \tau^s = \omega^{rs} \tau^s \sigma^r \,, \quad \omega = e^{2\pi i/N} \,, \quad r, s \in \mathbb{Z}_N \,, \tag{2.6}$$

and as obvious from the definition, σ_i, τ_j commute if $i \neq j$. Note that for $N = 2$: $\sigma = \sigma^z$, $\tau = \sigma^x$. The space of linear operators on $\mathcal{H}$ is spanned by monomials $\prod_{i=1}^L \sigma_i^{p_i} \tau_i^{q_i}$, $p_i, q_i \in \mathbb{Z}_N$.

The global symmetry $G = \mathbb{Z}_N$ is generated by the operator shifting the labels by one unit on every site:

$$P = \prod_{i=1}^L \tau_i^\dagger \,, \tag{2.7}$$

and each tensorand $\mathbb{C}^N$ in the Hilbert space $\mathcal{H}$ is direct sum of N one-dimensional irreducible representations of $\mathbb{Z}_N$. The Hamiltonians H we are going to consider will be invariant under the action of $\mathbb{Z}_N$. In other words, $[H, P] = 0$.

---

[2] These are the relations of the Heisenberg group which is defined by exponentiation of the Weyl algebra $[Q, P] = QP - PQ = i\mathbf{1}$. In terms of the generators $\sigma = \exp(iP)$, $\tau = \exp(iQ)$ and the central element $Z = \exp(i\mathbf{1})$ the group multiplication reduces to $\sigma^r \tau^s = Z^{rs} \tau^s \sigma^r$ for any parameters $r, s \in \mathbb{R}$.

2.2 Jordan Wigner transformation

Now we perform a generalized Jordan Wigner transformation to define the parafermions [23]. First we define operators living on the dual lattice as:

$$\mu_{i+1/2} = \sigma_i \sigma^\dagger_{i+1}\,, \quad \rho_{i+1/2} = \prod_{k=1}^{i} \tau^\dagger_k\,, \tag{2.8}$$

so that $\tau_i = \rho_{i-1/2}\rho^\dagger_{i+1/2}$, and μ, ρ satisfy the same commutation relations as respectively τ, σ. ρ is the disorder operator dual to the order operator σ, implementing the action of symmetries on spins between sites 1 and i, namely shifting those spins by one unit. Parafermions are then defined on middle points between vertices of the original and dual lattice, as the product of order and disorder variables:

$$\gamma_{2i-1} = \left(\prod_{k=1}^{i-1} \tau_k\right) \sigma_i\,, \quad \gamma_{2i} = \omega^{(N-1)/2} \left(\prod_{k=1}^{i-1} \tau_k\right) \sigma_i \tau_i\,. \tag{2.9}$$

They satisfy the following commutation relations

$$\gamma_i^N = 1\,, \quad \gamma_i \gamma_j = \omega \gamma_j \gamma_i \quad (\text{for } 1 \leq i < j \leq 2L)\,, \tag{2.10}$$

and the definition can be inverted to yield:

$$\sigma_i \sigma^\dagger_{i+1} = \omega^{(N+1)/2} \gamma_{2i} \gamma^{-1}_{2i+1} \tag{2.11}$$

$$\tau_i = \omega^{-(N-1)/2} \gamma^{-1}_{2i-1} \gamma_{2i}\,. \tag{2.12}$$

γ's are the generators of an algebra which generalize the Clifford algebra of "Majorana fermions" occurring at $N = 2$ [28]. Elements of a generalized Clifford algebra (with even number of generators, as in the case at hand, where the algebra is isomorphic to the complex matrices acting on the Hilbert space of the chain) are the sum of homogeneous elements X which have a well defined $\mathbb{Z}_N$-grading $|X| \in \mathbb{Z}_N$ given by:

$$PXP^{-1} = \omega^{|X|} X\,, \tag{2.13}$$

with P the symmetry generator expressed in terms of the parafermions as

$$P = \prod_{i=1}^{L} \omega^{(N-1)/2} \gamma_{2i} \gamma^{-1}_{2i-1}\,. \tag{2.14}$$

In particular, the parafermionic generators themselves are elements of degree 1.

3 Symmetry broken phases in $\mathbb{Z}_N$ spin chains

The general theory developed in [27, 29] classifies gapped phases of spin chains with symmetry group G in terms of classes of projective representations of subgroups H to which the symmetry is spontaneously broken. The subgroups of $\mathbb{Z}_N$ are given by $\mathbb{Z}_n$, where n is a divisor of N. $\mathbb{Z}_N$ only has trivial projective representations for any N (see [30, Appendix J]), so that gapped phases of $\mathbb{Z}_N$ spin chains are indexed by divisors of N and fully characterized by spontaneous symmetry breaking (SSB).

Statistical systems with $\mathbb{Z}_N$ symmetry have been studied extensively over the years, and their phase diagrams have been discussed in a variety of places, see e.g. [31, 32, 24, 33]. In this section, we explicitly construct representatives of all gapped phases we expect to have. On the way, we shall emphasize important aspects related to order parameters which will play a key role in the description of parafermionic phases in Section 4.

3.1 Construction of the phases

We first briefly recall the defining features of SSB in $\mathbb{Z}_N$ spin chains. The breaking of $\mathbb{Z}_2$ symmetry in the ordered phase of the quantum Ising chain is an elementary textbook example [34]: the phase is characterized by a non-vanishing expectation value of the order parameter σ. The generalization to the breaking of $\mathbb{Z}_N$ down to $\mathbb{Z}_n$, n divisor of N, is straightforward.

In the thermodynamic limit, the local order parameters describing the properties of the spin chain are given by the powers $\sigma^{\alpha n}$ with $\alpha \in \mathbb{Z}_m - \{0\}$ and $m = N/n$. The residual symmetry which leaves the order parameter invariant is $\mathbb{Z}_n$. It is generated by P^m, the m^{th} power of the global parity. (If $n = N$ the phase is disordered.) Elements of $\mathbb{Z}_N/\mathbb{Z}_n = \mathbb{Z}_m$ are generated by P^n, and permute the ground states cyclically among themselves, so that the ground state manifold is m–fold degenerate. We will refer to $\mathbb{Z}_m$ as the broken symmetries. The degeneracy of the ground state encodes a permutation representation of the symmetries on the ground state manifold. This representation corresponds to an invariant of the phase and cannot change unless a phase transition occurs [27]. The symmetry breaking perturbation $h \sum_i (\sigma_i^{\beta n} + \text{h.c.})$ (h small and positive) of the Hamiltonian will select a single ordered ground state, leading to a non-zero expectation value of the order parameter even when the external field h is set to zero.

For later convenience we will now explicitly construct representatives of each phase. This includes both the desired states and an associated parent Hamiltonian. According to the above discussion, a phase labeled by n should have m ordered ground states $|\psi_\alpha\rangle$ ($\alpha \in \mathbb{Z}_m$) which satisfy

$$P|\psi_\alpha\rangle = |\psi_{\alpha+1}\rangle\,, \quad P^{km}|\psi_\alpha\rangle = |\psi_\alpha\rangle\,, \tag{3.1}$$

for all $k \in \mathbb{Z}_n$. A simple solution of these constraints is given by the product states

$$|\psi_\alpha\rangle = \bigotimes_{i=1}^{L} |\alpha\rangle\!\rangle_i\,, \qquad \text{where} \qquad |\alpha\rangle\!\rangle := \sum_{k=0}^{n-1} |\alpha + mk\rangle\,. \tag{3.2}$$

The states $|\alpha\rangle\!\rangle$ are orbits under the subgroup $\mathbb{Z}_n$ of $\mathbb{Z}_N$. As such they span the subspace $\mathbb{C}^m \subseteq \mathbb{C}^N$ and they satisfy the periodicity condition $|\alpha\rangle\!\rangle \equiv |\alpha + m\rangle\!\rangle$.

The associated Hamiltonian will be defined in such a way that the m-fold degeneracy of the ground state is already exact in finite systems and not only in the thermodynamic limit. This is achieved by choosing a frustration free combination of specific projectors which are localized on two neighboring sites. Taking $H = -\sum_i h_{i,i+1}$, the following choice of the two-body term $h_{i,i+1}$ guarantees the $\mathbb{Z}_N$-invariance and the projector property ($h_{i,i+1}^2 = h_{i,i+1}$, $h_{i,i+1}^\dagger = h_{i,i+1}$)

$$h_{i,i+1} = \frac{1}{n^2} \sum_{\alpha=0}^{m-1} |\alpha\rangle\!\rangle_i |\alpha\rangle\!\rangle_{i+1} \langle\!\langle\alpha|_i \langle\!\langle\alpha|_{i+1}\,. \tag{3.3}$$

The Hamiltonian H is a sum of commuting projectors, so that it is possible to diagonalize all terms simultaneously. The resulting spectrum is given by the integers between $-L$ and 0, and hence the chain is gapped as desired. Assuming periodic or open boundary conditions will make no difference for the conclusions we want to draw in this section. In both cases the space of ground states is spanned by the vectors $|\psi_\alpha\rangle$ with $\alpha \in \mathbb{Z}_m$, for which $h_{i,i+1} = 1$.

It remains to find a more explicit form of the Hamiltonian (3.3) in terms of the spin operators σ_i and τ_i. Using the single-site matrix elements

$$\frac{1}{n}\langle q|\alpha\rangle\!\rangle\langle\!\langle\alpha|p\rangle = \frac{1}{N}\langle q| \sum_{\beta=0}^{m-1} (\sigma\omega^{-\alpha})^{n\beta} \sum_{j=0}^{n-1} \tau^{mj} |p\rangle\,, \tag{3.4}$$

it can easily be verified that Eq. (3.3) reduces to

$$h_{i,i+1} = \frac{m}{N^2} \sum_{\beta=0}^{m-1} (\sigma_i \sigma_{i+1}^\dagger)^{\beta n} \sum_{j,k=0}^{n-1} \tau_i^{jm} \tau_{i+1}^{km} \,. \tag{3.5}$$

We have thus succeeded in constructing gapped Hamiltonians realizing each possible SSB phase of $\mathbb{Z}_N$ spin chains.

Before moving on, we would like to comment on the specific form of the Hamiltonians obtained. The cases $n = 1$ and $n = N$ correspond, respectively, to the situations with no residual symmetry and full symmetry of the ground state. In these two extreme cases, the interaction simplifies to the expressions

$$(n, m) = (1, N) : \quad h_{i,i+1} = \frac{1}{N} \sum_{\alpha=0}^{N-1} (\sigma_i \sigma_{i+1}^\dagger)^\alpha \tag{3.6}$$

$$(n, m) = (N, 1) : \quad h_{i,i+1} = \frac{1}{N^2} \sum_{j,k=0}^{N-1} \tau_i^j \tau_{i+1}^k \,. \tag{3.7}$$

It is now evident that the general formula (3.5) is obtained by multiplying together the Hamiltonians of a $\mathbb{Z}_m$ chain whose symmetry is completely broken, $1/m \sum_{\beta=0}^{m-1} (\sigma_i^n (\sigma_{i+1}^\dagger)^n)^\beta$, and that of a $\mathbb{Z}_n$ chain without symmetry breaking, $1/n^2 \sum_{j,k=0}^{n-1} (\tau_i^m)^j (\tau_{i+1}^m)^k$.

While our definition reproduces the conventional form of the ferromagnetic clock model deep in the symmetry broken phase [18] for $n = 1$, the case $n = N$ leads to a more complicated expression than the one used in [18], namely $\sum_{j=0}^{N-1} \tau_i^j$ (even though both Hamiltonians have the same symmetric ground state). Moreover, the duality transformations (2.8) between the ordered and the disordered phase do not preserve the structure of the Hamiltonians (3.5). It is therefore reasonable to search for a slight modification of the Hamiltonian (3.5) such that it reduces to the form of the spin chains considered in [18], while retaining the ground states above. This goal is achieved by considering the expression

$$\hat{H} = -\sum_i \left(\frac{1}{m} \sum_{\beta=0}^{m-1} (\sigma_i \sigma_{i+1}^\dagger)^{\beta n} + \frac{1}{n} \sum_{j=0}^{n-1} \tau_i^{jm} \right) \,. \tag{3.8}$$

This operator is Hermitian, $\mathbb{Z}_N$ invariant, and each single term is a projector commuting with all the other terms. Therefore the chain is still gapped after the modification, albeit with a different spectrum in comparison to H.

The ground states are such that each term is 1 and $\hat{H}$ has exactly m ground states which are given by (3.2). Indeed, the terms $1/n \sum_{j=0}^{n-1} \tau_i^{jm}$ act as identity on states $\bigotimes_{i=1}^L |\alpha_i\rangle\!\rangle_i$ ($\alpha_i \in \mathbb{Z}_m$). Demanding, in addition, that also $1/m \sum_{\beta=0}^{m-1} (\sigma_i \sigma_{i+1}^\dagger)^{\beta n}$ acts as identity selects states with $\alpha_i = \alpha_{i+1}$. Hence $\hat{H}$ is another representative of the SSB phase under consideration. Note that the Hamiltonians $\hat{H}$ in (3.8) admit a simple action (in the thermodynamics limit) under the duality transformations (2.8). The order-disorder duality simply swaps the phases labeled by n with those labeled by m, and at the same time interchanges σ with ρ and τ with μ. Nonetheless, we will stick to our original definition of the Hamiltonian, see Eq. (3.5), in what follows.

3.2 Order parameters

In a macroscopic system with broken symmetry one will not observe a quantum superposition of ground states. Fluctuations will rather single out a unique ground state with a well-defined magnetization. In order to select a single ground state $|\psi_0\rangle$, we add a symmetry breaking perturbation $h \sum_i (\sigma_i^{\beta n} + \text{h.c.})$

to the Hamiltonian (3.5). Afterwards we take the thermodynamic limit and set $h = 0$. It is easy to check that this procedure implies a spontaneous magnetization

$$\langle \sigma_i^{\beta n} \rangle := \langle \psi_0 | \sigma_i^{\beta n} | \psi_0 \rangle \neq 0 \,, \tag{3.9}$$

and also $\langle \tau_i^{km} \rangle \neq 0$ for all $k \in \mathbb{Z}_n$. More importantly, we also have

$$\langle \rho_{i+1/2}^{km} \rangle \neq 0 \,. \tag{3.10}$$

In contrast, all other powers of σ and ρ have vanishing expectation values. Now we expect that slight modifications of Hamiltonians of gapped phases cannot change abruptly the expectation values unless a phase transition is reached. Then Eqs. (3.9, 3.10) hold not only for the finely tuned Hamiltonians we have constructed but in a whole region of non-zero measure in the phase diagram, which constitutes the SSB phase.

The mechanism of symmetry breaking is well understood for the Ising chain and it applies similarly to $\mathbb{Z}_N$ spins. The operator $\rho_{i+1/2}$ disorders an ordered state by creating kink excitations, topological objects rotating all spins from site 1 to i by one unit. The disordered phase can be characterized by the condensation of these kinks [35].

We are now prepared to appreciate that a particular situation is encountered when there exist two integers α, k such that $\alpha n = km$. Indeed, under these circumstances we are able to construct another *non-local* order parameter

$$(\rho_{i-1/2}^{\dagger} \sigma_i)^{\alpha n} \,. \tag{3.11}$$

(The non-zero expectation value is implied by Eqs. (3.9, 3.10).) With the definition $g := \gcd(n, m)$, the solutions of the equation $\alpha n = km$ can be parametrized as follows,

$$\alpha = k \frac{m}{n} \qquad \text{with} \qquad k \in \left\{0, \tfrac{n}{g}, 2\tfrac{n}{g}, \ldots, (g-1)\tfrac{n}{g}\right\} = \tfrac{n}{g}\mathbb{Z}_g \,. \tag{3.12}$$

We conclude that we have additional (non-local) order parameters whenever $g > 1$. Their existence is a direct consequence of being in a SSB phase of the $\mathbb{Z}_N$ spin chains. (Of course the presence of non-local order parameters does not mean that the phase at hand is topological, since the degeneracy of the ground state of the spin chain can be completely lifted through SSB.) In Section 4.2, we will shed more light onto the physical implications of the presence of this new non-local order parameter. It is no coincidence that the product (3.11) of order and disorder variables that we have just encountered agrees with powers of the parafermion, see Eq. (2.9).

4 Classification of parafermionic phases

In this section we will translate the classification of symmetry breaking patterns in $\mathbb{Z}_N$ spin chains into the dual language of parafermions. It will be found that the resulting parafermionic Hamiltonians generally exhibit a mixture of symmetry breaking and non-trivial topology which can be made manifest in the ground state degeneracies and in the occurrence of protected edge modes. We shall point out certain subtleties that arise when considering closed boundary conditions as opposed to open ones.

4.1 Duality transformation of the Hamiltonian

Our first goal is to rewrite the Hamiltonian Eq. (3.5) in terms of the parafermions. Using the transformation rules (2.11) and the relation

$$(\gamma_i^a \gamma_j^b)^c = \omega^{-abc(c-1)/2} \gamma_i^{ac} \gamma_j^{bc} \quad (\text{for } i < j) \,, \tag{4.1}$$

we immediately obtain the two-site Hamiltonian

$$h_{i,i+1} = \frac{m}{N^2} \sum_{\beta=0}^{m-1} \omega^{\beta n(N+\beta n)/2} \gamma_{2i}^{\beta n} \gamma_{2i+1}^{-\beta n} \times \sum_{j,k=0}^{n-1} \omega^{(m^2(j^2+k^2)-Nm(j+k))/2} \gamma_{2i-1}^{-jm} \gamma_{2i}^{jm} \gamma_{2i+1}^{-km} \gamma_{2i+2}^{km} \,. \tag{4.2}$$

Despite the non-locality of the transformation, the Hamiltonian still remains local when expressed in terms of the parafermionic variables. The only complication arises for periodic boundary conditions since the naive application of the Jordan Wigner transformation generates a non-local term in this case. In other words, the correct Hamiltonian for the periodic parafermionic chain needs to be different from that of the periodic spin model, see Section 4.3.2 for details.

In the following we will use a basis of parity eigenvectors in the space of ground states which is more natural to work with in the parafermionic picture:

$$|\phi_\alpha\rangle := \sum_{\beta=0}^{m-1} \omega^{\beta\alpha n} |\psi_\beta\rangle \,, \tag{4.3}$$

so that

$$\sigma_i^{\beta n} |\phi_\alpha\rangle = |\phi_{\alpha+\beta}\rangle \,, \quad \tau_i^{km} |\phi_\alpha\rangle = |\phi_\alpha\rangle \,, \quad P|\phi_\alpha\rangle = \omega^{\alpha n} |\phi_\alpha\rangle \,. \tag{4.4}$$

4.2 Quantum phases

In Section 3 we characterized SSB phases of $\mathbb{Z}_N$ spin chains. However, a moment of thought reveals that the local order parameter $\sigma^{\alpha n}$ of the spin chain becomes non-local when expressed in terms of parafermionic variables. Since physical perturbations are assumed to be local, this observation suggests that there is no way to spontaneously break the symmetry in the parafermionic chain, and that the phases are characterized by topological order instead of ferromagnetic order. In this case, the states emerging at the macroscopic level would be those with definite $\mathbb{Z}_N$ parity. In the fermionic case $N = 2$, for example, it is well-established that there is no local order parameter, and correspondingly no local perturbation which can break the symmetry.

While valid for a considerable subset of symmetry breaking patterns, the previous expectation fails to be true in general. This exactly happens in the phases labeled by a divisor n such that $g = \gcd(n, m) > 1$. Parafermionic parity can be broken in these cases and, depending on the precise values of g and n, this leads to phases which, in addition to ferromagnetic order, might as well exhibit topological order.

In order to substantiate our claim, let us briefly recall our results from Section 3.2. For $g > 1$ we have pointed out the existence of a second set of order parameters (3.11), which when rewritten in terms of parafermions become local. More precisely, they correspond exactly to mutually commuting powers of the parafermions,

$$\left[\gamma_i^{pN/g}, \gamma_j^{p'N/g}\right] = 0 \quad (\text{with } p, p' \in \mathbb{Z}_g) \,. \tag{4.5}$$

Note that, more generally, $\gamma_j^{pN/g}$ commutes with $\gamma_i^{\alpha n}$ and γ_i^{km} for every i, j, so that:

$$\left[\gamma_j^{pN/g}, h_{i,i+1}\right] = 0 \,. \tag{4.6}$$

In order to understand the physical implications of the order parameters let us then consider a $\mathbb{Z}_N \to \mathbb{Z}_N/\mathbb{Z}_g$ symmetry breaking perturbation of the original parafermionic Hamiltonian,

$$H \;\mapsto\; H + h \sum_i \left\{ \gamma_i^{pN/g} + \text{h.c.} \right\} \quad (\text{with } p \in \mathbb{Z}_g - \{0\}) \,. \tag{4.7}$$

We suppose open boundary conditions, for which the Hamiltonians for the spins and the parafermions are identical. The extra term in the Hamiltonian induces the following action on the ground states,

$$\gamma_{2i-1}^{pN/g}|\psi_\alpha\rangle = \left(\prod_{k=1}^{i-1}\tau_k\right)^{pN/g} \sigma_i^{pN/g}|\psi_\alpha\rangle = \omega^{\alpha pN/g}|\psi_\alpha\rangle\,, \tag{4.8}$$

and analogously for γ_{2i}, with an extra α-independent phase. As a consequence, the SSB induced by the perturbation (4.7) singles out a subset of the m ground states, exactly those labeled by an element α such that

$$\alpha\frac{pN}{g} = 0 \pmod{N}\,. \tag{4.9}$$

It is evident that this equation has $\tilde{m} = m/g$ solutions

$$\alpha \in \{0, g, \ldots, g\tilde{m}-1\} = g\mathbb{Z}_{\tilde{m}}\,. \tag{4.10}$$

We believe that the degeneracy determined by the number of the remaining ground states cannot be lifted by SSB, and should therefore be regarded as topological. (If however $m = g$ no such degeneracy is present, as e.g. when $n = 2, N = 4$, and the system is in a purely symmetry breaking phase.)

Our previous interpretation receives additional support when considering the system with periodic boundary conditions. Namely, in that case the ground state degeneracy due to topologically protected edge modes is expected to disappear while the degeneracy due to symmetry breaking should persist. The degeneracy due to symmetry breaking is g, and therefore we expect a closed parafermionic chain to have a g-fold degenerate ground state. Whenever $g < m$ our arguments implies the existence of topologically protected edge modes. We illustrate this phenomenology in the case of $\mathbb{Z}_8$ parafermions in figure 1. Our expectations will be verified in Section 4.3 using the explicit realization of the phases.



Figure 1: (Color on-line) Degeneracies of the ground states as a function of the divisor n labeling the phases of $\mathbb{Z}_8$ parafermions for both open (left) and closed (right) boundary conditions.

4.3 Realization of the phases

In this section we will provide an explicit realization of all parafermionic phases discussed in the previous section, thereby putting our considerations on a concrete basis. It turns out to be convenient to discuss open and closed boundary conditions separately.

4.3.1 Open boundaries

The Hamiltonian for the open chain is

$$H_{\text{op}} = -\sum_{i=1}^{L-1} h_{i,i+1}\,, \tag{4.11}$$

where $h_{i,i+1}$ is given in (4.2). H_{op} has m degenerate ground states $|\phi_\alpha\rangle, \alpha \in \mathbb{Z}_m$, see Eq. (4.3). When the phases are purely topological, namely when $g := \gcd(n, m) = 1$, the degeneracy can be understood in terms of zero modes (operators commuting with Hamiltonian H_{op}) γ_1^n and γ_{2L}^n which are localized at the boundary. Indeed, acting with powers of γ_1^n (or γ_{2L}^n) on a given ground state, say $|\phi_0\rangle$, produces the other $m-1$ ground states distinguished by parity eigenvalues. In the generic case when $g \geq 1$, the ground state degeneracy is still due to zero modes, but now some of the m zero modes are no longer localized at the boundary. From Eq. (4.6) we see that $\left[\gamma_j^{pN/g}, H_{\mathrm{op}}\right] = 0$ for every j, and acting with $\gamma_j^{pN/g}$ on $|\phi_0\rangle$ will produce a state proportional to $|\phi_{pm/g}\rangle$, so that these zero modes are not located at the edge. These zero modes are precisely those responsible for the degeneracy due to symmetry breaking. In the particular case of purely topological phase labeled by $n = 1$ with $m = N$ degenerate ground states, the question of robustness of the edge modes away from the finitely tuned Hamiltonians was raised and answered in [18]. We expect that all the zero modes responsible for topological degeneracy we found will also stay localized at the boundary in the whole phase.

It has been pointed out in [6, 36] that the quantum phases not distinguished by symmetry breaking (pure topological phases) correspond to different ways to "fractionalize" the symmetry operators on the space $\mathcal{H}_{\mathrm{low}}$ of low energy states of the open chain (or low entanglement energy states of the closed chain). According to these ideas, parity should be represented as the operator $\hat{P} = P_l P_r$ on $\mathcal{H}_{\mathrm{low}}$ in our case, with P_l and P_r acting non-trivially only at the left and right boundaries, respectively. Denoting by x the $\mathbb{Z}_N$ degree of P_l (and by $-x$ that of P_r), consistency of the factorization requires that [36]

$$\hat{P} P_l = \omega^x P_l \hat{P} = P_l P_r P_l = \omega^{x^2} P_l \hat{P} \,. \tag{4.12}$$

In other words, the degree of P_l must satisfy the equation

$$x(x-1) = 0 \pmod{N} \,. \tag{4.13}$$

The number of solutions to this equation with $x \in \mathbb{Z}_N$ is given by the number of divisors n of N such that n and N/n are co-prime (see below).

In what follows we will connect our previous considerations from Section 4.2 to the prediction obtained from symmetry fractionalization. Equation (4.13) selects out of the possible phases a distinguished subset for which a low energy representation of parity exists. In those cases, the fractionalized symmetry operator can be constructed explicitly for our Hamiltonians (4.11), for which $\mathcal{H}_{\mathrm{low}}$ coincides with the ground state manifold. Since the edge modes are low energy operators localized at the boundary we make the ansatz

$$\hat{P} = \omega^{r(N+r)/2} \gamma_1^r \gamma_{2L}^{N-r} \,. \tag{4.14}$$

The operator $\hat{P}$ satisfies the defining property of parity $\hat{P}^N = 1$, is of degree zero, and it acts on the space of ground states as

$$\hat{P}|\phi_\alpha\rangle = \sigma_1^r (\sigma_L^\dagger)^r P^r |\phi_\alpha\rangle = \omega^{\alpha n r} \sum_{\beta=0}^{m-1} \omega^{\beta \alpha n} \sigma_1^r |\beta\rangle\!\rangle \otimes \cdots \otimes (\sigma_L^\dagger)^r |\beta\rangle\!\rangle \tag{4.15}$$

$$= \omega^{\alpha n r} \sum_{\beta=0}^{m-1} \omega^{\beta \alpha n} \sum_{k=0}^{n-1} \omega^{(\beta+mk)r} |\beta + mk\rangle \otimes \cdots \otimes \sum_{j=0}^{n-1} \omega^{-(\beta+mj)r} |\beta + mj\rangle \,. \tag{4.16}$$

If we demand that $\hat{P}$ acts as P on the space of ground states, this requires that $mkr = 0 \pmod{N}$ for every k, that is $r = n\epsilon$, $\epsilon \in \mathbb{Z}_m$. In this case, we have:

$$\hat{P}|\phi_\alpha\rangle = \omega^{\alpha n^2 \epsilon} |\phi_\alpha\rangle \,, \tag{4.17}$$

and ϵ must be determined by the equation $\alpha n(n\epsilon - 1) = 0 \pmod N$ for every $\alpha \in \mathbb{Z}_m$. Clearly, the case $\alpha = 1$ implies the others, so that we have to find an ϵ such that

$$n(n\epsilon - 1) = 0 \pmod N\,. \tag{4.18}$$

Alternatively, we can rewrite the previous equation as

$$n\epsilon = 1 + pm \quad \text{with} \quad p \in \mathbb{Z}\,. \tag{4.19}$$

This equation has no solution if $g := \gcd(n,m) > 1$. Indeed, in that case the left hand side of $(n\epsilon - pm)/g = 1/g$ is an integer while the right hand side is not. On the other hand, if $g = 1$ there exists a unique solution.[3] This result is consistent with Section 4.2, where purely topological phases were predicted to occur only if $g = 1$.

What is left to do is to show that we have exhausted all the purely topological phases predicted by Eq. (4.13). The operator $\hat{P}$ we have constructed, is obviously of the form $\hat{P} = P_l P_r$ with degree of P_l equal to $n\epsilon$, and Eq. (4.13) is clearly implied by (4.18). We now prove that, furthermore, every solution of $x(x-1) = 0 \pmod N$ defines a divisor n with $\gcd(n, N/n) = 1$, so that solutions of Eq. (4.13) and (4.18) are in bijection. The proof is easily carried out. Setting $n = \gcd(x,N)$, simple properties of gcd give:

$$\gcd(n, N/n) = \gcd\left(\gcd(x,N), \frac{N}{\gcd(x,N)}\right) = \frac{\gcd(\gcd(x,N)^2, N)}{\gcd(x,N)} \tag{4.20}$$

$$= \frac{\gcd(\gcd(x^2,N^2),N)}{\gcd(x,N)} = \frac{\gcd(x^2, \gcd(N^2,N))}{\gcd(x,N)} \tag{4.21}$$

$$= \frac{\gcd(x^2 \pmod N, N)}{\gcd(x,N)} = \frac{\gcd(x,N)}{\gcd(x,N)} = 1\,. \tag{4.22}$$

Therefore all the solutions of (4.13) can be expressed by those of (4.18) in the form $x = n\epsilon$.

We conclude with some more technical notes. Divisors n of N such that $\gcd(n, N/n) = 1$ are called unitary. In the prime factorization $N = \prod p_i^{a_i}$, $n = \prod p_i^{c_i}$ is a unitary divisor if each c_i is 0 or a_i. For example the divisors of $N = 4$ are $1, 2, 4$ but only $1, 4$ are unitary. The number of parafermionic phases characterized by effective parity (exhibiting then pure topological order) is given by the number of unitary divisors. For $N = 1, 2, 3, \ldots$ their numbers are

$$1,2,2,2,2,4,2,2,2,4,2,4,2,4,4,2,2,4,2,4,4,4,2,4,2,4,2,4,2,8,\ldots \quad \text{(Sloane's A034444)}\,.$$

These numbers can also be identified with 2^q, where q is the number of different primes dividing N.

4.3.2 Closed boundaries

The parafermionic closed chain is defined by the local Hamiltonian

$$H^P_{\text{per}} = H_{\text{op}} - h^P_{L,1}\,. \tag{4.23}$$

In specifying $h^P_{L,1}$ we need to pay attention to the order of indices since parafermions on different sites do not commute. The substitution $(2i, 2i+1) \to (2L, 1)$ in the term $\gamma_{2i}^{\beta n}\gamma_{2i+1}^{-\beta n}$ of Eq. (4.2) inverts the order of indices, and two different Hamiltonians are obtained depending on whether we commute the parafermions before or after the replacement of indices. Indeed performing the above substitution and then commuting the parafermions gives $\gamma_{2L}^{\beta n}\gamma_1^{-\beta n} = \omega^{(\beta n)^2}\gamma_1^{-\beta n}\gamma_{2L}^{\beta n}$. If instead we commute the parafermions in Eq. (4.2) before replacing the indices we will get $\omega^{-(\beta n)^2}\gamma_1^{-\beta n}\gamma_{2L}^{\beta n}$. The ambiguous extra phase produced by inverting the operations signals an ambiguity in the definition of $h^P_{L,1}$. The correct definition of the coupling instead keeps the order of labels and implements the replacement $(2i, 2i+1) \to (1, 2L)$ in $\gamma_{2i}^{\beta n}\gamma_{2i+1}^{-\beta n}$. Following this prescription we get

---

[3] Uniqueness is easy to show since if $\epsilon_1 = (1 + p_1 m)/n$ and $\epsilon_2 = (1 + p_2 m)/n$ are two solutions, then $\epsilon_1 - \epsilon_2 = (p_1 - p_2)m/n$ must be an integer, implying $(p_1 - p_2)$ is a multiple of n and so $\epsilon_1 - \epsilon_2 = 0 \pmod m$.

$$h^P_{L,1} = \frac{m}{N^2}\sum_{\beta=0}^{m-1} \omega^{\beta n(N+\beta n)/2}\gamma_1^{\beta n}\gamma_{2L}^{-\beta n} \times \sum_{j,k=0}^{n-1} \omega^{(m^2(j^2+k^2)-Nm(j+k))/2}\gamma_{2L-1}^{-jm}\gamma_{2L}^{jm}\gamma_1^{-km}\gamma_2^{km}\,. \tag{4.24}$$

When rewritten in terms of spins, this operator turns out to be non-local. More precisely, it contains a twisting by the inverse global parity $P^\dagger$:

$$h^P_{L,1} = \frac{m}{N^2}\sum_{\beta=0}^{m-1}(P^\dagger\sigma_L\sigma_1^\dagger)^{\beta n}\sum_{j,k=0}^{n-1}\tau_L^{jm}\tau_1^{km}\,. \tag{4.25}$$

While completely equivalent when regarded with open boundary conditions, the spin model and the parafermionic chain thus exhibit essential differences when considered on a closed ring.

In order to understand the physical implications of our previous comment, we now study the spectrum of the periodic chain, with particular attention to the ground state and its degeneracy. Recall that H_{op} has m ground states which are given by $|\phi_\alpha\rangle$ as defined in Eq. (4.3). Since $[H_{\mathrm{op}}, h^P_{L,1}] = 0$, the operators H_{op} and $h^P_{L,1}$ admit a common basis of eigenvectors, and our problem simply amounts to diagonalizing the projector $h^P_{L,1}$ on the space spanned by the vectors $|\phi_\alpha\rangle$.

A brief calculation shows that the operator $h^P_{L,1}$ acts on the ground states of the open chain as

$$h^P_{L,1}|\phi_\alpha\rangle = \frac{1}{m}\sum_{\beta=0}^{m-1}\omega^{-\beta n^2\alpha}|\phi_\alpha\rangle = \delta_{\alpha n=0 \pmod{m}}|\phi_\alpha\rangle\,. \tag{4.26}$$

Put differently, the state $|\phi_\alpha\rangle$ remains in the ground state manifold of the closed parafermionic chain if and only if

$$\alpha n = 0 \pmod{m}\,. \tag{4.27}$$

In order to enumerate the solutions to this equation, let us first introduce the symbols $\tilde{n} = n/g$ and $\tilde{m} = m/g$ where we used the previous abbreviation $g = \gcd(n, m)$. With this notation, we can write the solutions of Eq. (4.27) as

$$\alpha = p\frac{\tilde{m}}{\tilde{n}} \quad \text{with} \quad p \in \tilde{n}\mathbb{Z}_g\,. \tag{4.28}$$

Hence, the possible values of p are $p = 0, \tilde{n}, \ldots, \tilde{n}(g-1)$, so that there are g distinct solutions.

It may easily be shown that the previous g states exhaust the ground space manifold of the periodic Hamiltonian (4.23). Note that we can associate the degeneracy of the closed chain to parafermionic zero modes as in the case of the open chain. Indeed Eq. (4.6) implies that $\left[\gamma_i^{pN/g}, H^P_{\mathrm{per}}\right] = 0$, and acting with powers of $\gamma_i^{N/g}$ on a reference ground state will produce g mutually orthogonal ground states with different parity eigenvalues.

Our analysis implies that the closed parafermionic chains have a unique ground state for n being a unitary divisor, while there exists a g-fold degenerate ground state when n fails to be a unitary divisor. In the former case, there is no spontaneous symmetry breaking and the ground state degeneracy in the open chain may be completely attributed to the non-trivial topology of the system. On the other hand, in the latter case, the ground state degeneracy is only partially lifted when making the transition from open to closed boundary conditions. In that case, we observe a combination of non-trivial topology and spontaneous symmetry breaking if $m/g > 1$. For $g = m$ there is no non-trivial topology and only symmetry breaking. These results are in complete agreement with our expectations from Section 4.2.

5 Conclusions and outlook

In the current paper we have classified the massive phases of $\mathbb{Z}_N$ parafermionic chains. This was achieved by constructing all gapped symmetry broken phases in the dual quantum spin chains. The number of distinct phases and their specific characteristics depend crucially on the number–theoretic properties of N. First of all, it is easy to construct one SSB phase of the spin model for each divisor n of N. These phases are characterized by the (co-)existence of local and non-local order parameters. Since $\mathbb{Z}_N$ does not lead to symmetry protected topological phases [3, 29], the enumeration of divisors exhausts all possibilities for gapped phases.

In a second step we have then interpreted the resulting ground states and Hamiltonians from a parafermionic perspective. Of course, this leads to the same number of phases. However, in contrast to the spin model the parafermionic phases can now exhibit features of both, symmetry breaking and non-trivial topology. Whether the resulting parafermionic phase should be interpreted as topological first of all depends on whether n and N/n still have common divisors. A parafermionic phase is purely topological if n is a unitary divisor of N. By definition, this is the case if n and the quotient N/n are co-prime or, in our notation, if $g = \gcd(n, N/n) = 1$. The number of unitary divisors is known to be 2^q where q is the number of distinct prime factors in N. In contrast, a parafermionic phase described by $n < N$ not being a unitary divisor (i.e. $g > 1$) definitively exhibits spontaneous symmetry breaking. If, in addition, $gn < N$ then the phase is also protected by topology. Topology protected phases in open systems exhibit gapless parafermionic edge modes.

While our current results are only concerned with the intrinsic $\mathbb{Z}_N$ symmetry of parafermionic chains, it would be interesting to extend our analysis to systems which are required to respect additional symmetries such as inversion or time-reversal symmetry. Alternatively, one could also add internal degrees of freedom which transform under a discrete or continuous symmetry group. In all these cases one is lead to so-called symmetry protected topological phases which, for bosonic systems, have been fully classified in Refs. [3, 27, 29]. It should be noted that models with continuous symmetry lead to a few peculiarities that have been addressed in Ref. [37].

For fermionic chains it is known that the inclusion of additional symmetries has profound consequences. Most importantly, the restriction to time-reversal invariant systems enhances the group of topological invariants to $\mathbb{Z}$ or $\mathbb{Z}_8$, respectively, depending on whether interactions are allowed [38, 26, 6] or not [4, 5]. A general mathematical formalism for treating fermionic symmetry protected topological systems has been developed in Ref. [39] and it appears feasible to extend it to the $\mathbb{Z}_N$-graded generalization of the Clifford algebra [28] needed to describe parafermions.

Finally, the classification scheme used in this paper is, of course, not limited to $\mathbb{Z}_N$ clock models. It should rather be applicable to any model of statistical physics which exhibits a Kramers-Wannier type of duality. While abelian dualities are well-explored (see, e.g., [40]) and close to our exposition, recent progress on non-abelian dualities [41] might offer a vast playground for the exploration of new topological phases.

Needless to say, knowledge about the classification of topological phases, symmetry protected or not, and a representative Hamiltonian for each phase should merely be the starting point for a more thorough investigation. Indeed, the ultimate goal should be to gain control over the complete phases diagram of physically realistic Hamiltonians, including the location and the nature of phase transitions. For the Majorana chain with interactions such an analysis was provided in [42], based on the known phase diagram of the dual anisotropic next-nearest-neighbor Ising (ANNNI) model. It would be interesting to extend this analysis to the case of $\mathbb{Z}_N$ spin models. Also, with regard to the potential physical realization and detection of parafermions one will need to understand the effects of adding disorder to the couplings.

*** *Note* ***

While in the process of completing this work, the paper [36] appeared which has considerable overlap with our own results. We gratefully acknowledge useful discussions with J. Motruk and A. Turner about aspects of entanglement and symmetry fractionalization in parafermionic chains prior to the publication of their paper.

Acknowledgments

We would like to thank A. Altland for interesting discussions leading to the launch of this project and K. Duivenvoorden for collaboration in the initial stages and extremely useful discussions. Both authors are funded by the German Research Council (DFG) through M. Zirnbauer's Leibniz Prize, DFG grant no. ZI 513/2-1. Additional support is received from the DFG through the SFB|TR 12 "Symmetries and Universality in Mesoscopic Systems".

References

[1] A. Y. Kitaev, "Quantum computations: algorithms and error correction," *Russ. Math. Surv. 52* (1997) 1191–1249.

[2] M. Freedman, M. Larsen, and Z. Wang, "A modular functor which is universal for quantum computation," *Commun. Math. Phys. 227* (2002) 605–622, `arXiv:quant-ph/0001108`.

[3] X. Chen, Z.-C. Gu, and X.-G. Wen, "Classification of gapped symmetric phases in one-dimensional spin systems," *Phys. Rev. B83* (2011) 035107, `arXiv:1008.3745`.

[4] S. Ryu, A. P. Schnyder, A. Furusaki, and A. W. W. Ludwig, "Topological insulators and superconductors: tenfold way and dimensional hierarchy," *New J. Phys. 12* (2010) 065010, `arXiv:0803.2786`.

[5] A. Kitaev, "Periodic table for topological insulators and superconductors," *AIP Conf. Proc. 1134* (2009) 22–30, `arXiv:0901.2686`.

[6] A. M. Turner, F. Pollmann, and E. Berg, "Topological phases of one-dimensional fermions: An entanglement point of view," *Phys. Rev. B83* (2011) 075102, `arXiv:1008.4346`.

[7] A. Y. Kitaev, "Unpaired Majorana fermions in quantum wires," *Phys. Usp. 44* (2001) 131, `arXiv:cond-mat/0010440`.

[8] J. Alicea, Y. Oreg, G. Refael, F. von Oppen, and M. P. A. Fisher, "Non-abelian statistics and topological quantum information processing in 1D wire networks," *Nature Phys. 7* (2011) 412–417, `arXiv:1006.4395`.

[9] R. M. Lutchyn, J. D. Sau, and S. Das Sarma, "Majorana fermions and a topological phase transition in semiconductor-superconductor heterostructures," *Phys. Rev. Lett. 105* (2010) 077001, `arXiv:1002.4033`.

[10] Y. Oreg, G. Refael, and F. von Oppen, "Helical liquids and Majorana bound states in quantum wires," *Phys. Rev. Lett. 105* (2010) 177002, `arXiv:1003.1145`.

[11] V. Mourik, K. Zuo, S. M. Frolov, S. R. Plissard, E. P. A. M. Bakkers, and L. P. Kouwenhoven, "Signatures of Majorana fermions in hybrid superconductor-semiconductor nanowire devices," *Science 336* (2012) 1003–, `arXiv:1204.2792`.

[12] A. Das, Y. Ronen, Y. Most, Y. Oreg, M. Heiblum, and H. Shtrikman, "Zero-bias peaks and splitting in an Al-InAs nanowire topological superconductor as a signature of Majorana fermions," *Nature Phys. 8* (2012) 887–895, `arXiv:1205.7073`.

[13] D. Bagrets and A. Altland, "Class D spectral peak in Majorana quantum wires," *Phys. Rev. Lett. 109* (2012) 227005, `arXiv:1206.0434`.

[14] F. Pientka, G. Kells, A. Romito, P. W. Brouwer, and F. von Oppen, "Enhanced zero-bias Majorana peak in the differential tunneling conductance of disordered multisubband quantum-wire/superconductor junctions," *Phys. Rev. Lett. 109* (2012) 227006, `arXiv:1206.0723`.

[15] P. Neven, D. Bagrets, and A. Altland, "Quasiclassical theory of disordered multi-channel Majorana quantum wires," `arXiv:1302.0747`.

[16] D. J. Clarke, J. Alicea, and K. Shtengel, "Exotic non-Abelian anyons from conventional fractional quantum Hall states," *Nature Comm. 4* (2013) 1348, `arXiv:1204.5479`.

[17] N. H. Lindner, E. Berg, G. Refael, and A. Stern, "Fractionalizing Majorana fermions: Non-Abelian statistics on the edges of Abelian quantum Hall states," *Phys. Rev. X2* (2012) 041002, `arXiv:1204.5733`.

[18] P. Fendley, "Parafermionic edge zero modes in Z_n-invariant spin chains," *J. Stat. Mech. 11* (2012) 20, `arXiv:1209.0472`.

[19] M. Cheng, "Superconducting proximity effect on the edge of fractional topological insulators," *Phys. Rev. B86* (2012) 195126, `arXiv:1204.6084`.

[20] A. Vaezi, "Fractional topological superconductor with fractionalized Majorana fermions," *Phys. Rev. B87* (2013) 035132, `arXiv:1204.6245`.

[21] H. S. Green, "A generalized method of field quantization," *Phys. Rev. 90* (1953) 270–273.

[22] O. W. Greenberg and A. M. L. Messiah, "Selection rules for parafields and the absence of para particles in nature," *Phys. Rev. 138* (1965) B1155–B1167.

[23] E. H. Fradkin and L. Kadanoff, "Disorder variables and parafermions in two-dimensional statistical mechanics," *Nucl. Phys. B170* (1980) 1–15.

[24] V. Fateev and A. Zamolodchikov, "Parafermionic currents in the two-dimensional conformal quantum field theory and selfdual critical points in Z_N invariant statistical systems," *Sov. Phys. JETP 62* (1985) 215–225.

[25] N. Read and E. Rezayi, "Beyond paired quantum Hall states: Parafermions and incompressible states in the first excited Landau level," *Phys. Rev. B59* (1999) 8084–8092, `cond-mat/9809384`.

[26] L. Fidkowski and A. Kitaev, "Topological phases of fermions in one dimension," *Phys. Rev. B83* (2011) 075103, `arXiv:1008.4138`.

[27] N. Schuch, D. Pérez-García, and I. Cirac, "Classifying quantum phases using matrix product states and PEPS," *Phys. Rev. B84* (2011) 165139, `arXiv:1010.3732`.

[28] A. Morris, "On a generalized Clifford algebra," *Quart. J. Math. 18* (1967) 7–12.

[29] X. Chen, Z.-C. Gu, and X.-G. Wen, "Complete classification of one-dimensional gapped quantum phases in interacting spin systems," *Phys. Rev. B84* (2011) 235128, `arXiv:1103.3323`.

[30] X. Chen, Z.-C. Gu, Z.-X. Liu, and X.-G. Wen, "Symmetry protected topological orders and the group cohomology of their symmetry group," `arXiv:1106.4772v6`. Note: The relevant statements are missing in the published version in Science 338 (2012) 1604-1606.

[31] S. Ostlund, "Incommensurate and commensurate phases in asymmetric clock models," *Phys. Rev. B24* (1981) 398–405.

[32] F. C. Alcaraz and R. Köberle, "The phases of two-dimensional spin and four-dimensional gauge systems with Z(N) symmetry," *J. Phys. A14* (1981) 1169.

[33] G. Ortiz, E. Cobanera, and Z. Nussinov, "Dualities and the phase diagram of the p-clock model," *Nucl. Phys. B854* (2012) 780 – 814.

[34] S. Sachdev, *Quantum Phase Transitions*. Cambridge University Press, 2001.

[35] E. Fradkin and L. Susskind, "Order and disorder in gauge systems and magnets," *Phys. Rev. D17* (1978) 2637–2658.

[36] J. Motruk, A. M. Turner, E. Berg, and F. Pollmann, "Topological phases in gapped edges of fractionalized systems," `arXiv:1303.2194`.

[37] K. Duivenvoorden and T. Quella, "On topological phases of spin chains," `arXiv:1206.2462`. (accepted for publication in Phys. Rev. B).

[38] L. Fidkowski and A. Kitaev, "Effects of interactions on the topological classification of free fermion systems," *Phys. Rev. B81* (2010) 134509, `arXiv:0904.2197`.

[39] Z.-C. Gu and X.-G. Wen, "Symmetry-protected topological orders for interacting fermions – fermionic topological non-linear sigma-models and a group super-cohomology theory," `arXiv:1201.2648`.

[40] R. Savit, "Duality in field theory and statistical systems," *Rev. Mod. Phys. 52* (1980) 453–488.

[41] E. Cobanera, G. Ortiz, and E. Knill, "The non-Abelian duality problem," `arXiv:1206.1367`.

[42] F. Hassler and D. Schuricht, "Strongly interacting Majorana modes in an array of Josephson junctions," *New J. Phys. 14* (2012) 125018, `arXiv:1206.2134`.